

Complete Medium

RPMI 1640 (10% FBS)

1. Product Overview

Complete Medium RPMI 1640 (10% FBS) is formulated with all essential nutrients required for adherent cell culture, enabling robust cell growth and proliferation without supplementation of additional basal components.

Employing advanced dynamic freeze-drying technology, this product lyophilizes a complex nutrient system comprising PRMI1640 basal culture medium, 10% fetal bovine serum (FBS), antibiotics (dual antibiotics), glutamine, and other essential additives into a homogeneous powder, while retaining the native structural integrity and biological activity of all constituents. It can be rapidly and conveniently reconstituted via gentle mixing and shaking, delivering sufficient energy and comprehensive nutritional support for highly metabolically active and fast-proliferating cells.

Supplemented with 10% FBS and 1% penicillin–streptomycin solution, and featuring a stable pH range of 7.00–7.40, this product is commonly used in research on immune cell function, in vitro culture of tumor cells, and preparation of monoclonal antibodies from hybridoma cells.

2. Core Ingredients and Functions

The table below lists the major components of this complete culture medium and describes the role of each:

Component	Function	Typical Concentration
Basal culture medium (RMI1640)	basic nutritional support and a stable growth environment for cells cultured in vitro.	89 %
Fetal bovine serum (FBS)	Supplementing growth factors, hormones, and adhesion factors promotes cell proliferation and adhesion.	10%
Penicillin-Streptomycin	Inhibit bacterial contamination (penicillin: 100 U/mL; streptomycin: 100 µg/mL).	1%
L-glutamine	Key substances for cellular energy metabolism.	300mg/L
D-glucose	It provides the energy and raw materials needed for metabolism in cells cultured in vitro .	2.0 g/L
Sodium bicarbonate	Maintaining pH stability of the culture medium	2g/L

Note: Actual component composition may vary by product variant. Refer to the Certificate of Analysis (CoA) for lot-specific information.

3. Storage Conditions

Storage Temperature	<i>2 – 8 °C (Refrigerate; do not freeze)</i>
Protection from Light	<i>Store in the dark or in amber/foil-wrapped containers to prevent photodegradation</i>
Shelf Life (Unopened)	<i>24 months from date of manufacture</i>
Shelf Life (Opened)	<i>Use within 3-4 weeks after opening; aliquot to minimize freeze-thaw or repeated warming</i>
Container	<i>Maintain in original sealed bottle or sterile, airtight container</i>

4. Preparations Before Use

4.1 Visual inspection

- Confirm that the lyophilized culture medium powder has not dissolved, the vial is undamaged and leak-free, and the liquid is clear without turbidity or sediment. If any abnormalities are found, contact the supplier immediately for a replacement.

4.2 Equilibration

- Warm the medium to 37 °C in a water bath or incubator (15–30 minutes) before use.
- Do not warm more medium than needed for immediate use; discard any unused warmed medium at the end of the working day.

4.3 Sterility

- All components of this medium have been sterile-filtered (0.2 µm) by the manufacturer.
- Always handle medium under aseptic conditions using a certified biosafety cabinet (BSC).
- Inspect the bottle for leaks, damaged seals, or signs of contamination before use.

5. Product Specifications

Product Name	<i>Complete Medium RPMI 1640 (10% FBS)</i>
Product Code	<i>HD6193-100; HD6193-500; HD6193A-100; HD6193A-500</i>
Volume	<i>100ml/bottle, 10*100ml/box; 500ml/bottle</i>

6. Instructions for Use

6.1 Temperature Equilibration:

- Remove the reagent bottle from the 2 – 8°C refrigerator before use, and let it stand for 10 minutes to equilibrate to room temperature.

6.2 Powder Settling:

- Press the top of the bottle cap to allow the powder inside to fall smoothly.

6.3 Dissolution & Mixing:

- Shake the bottle vigorously for 1 minute, and observe until the powder is completely dissolved.

6.4 Removal of Powder Cartridge:

- Remove the bottle cap, gently twist it to detach the powder cartridge, and discard it.

6.5 Ready for Use:

- The reagent is ready for normal use once fully dissolved, with a pH range of 7.00–7.40.

7. Reminder

- For Research Use Only (RUO). This product is not intended for use in diagnostic procedures or therapeutic applications.
- Under ambient temperature conditions, the product can maintain its quality for up to one week. It is recommended to use ice packs for transportation; **do not use dry ice**. In northern regions where freezing may occur in winter, please thaw the product completely before storage. Store the product in a refrigerator at 2–8°C.
- Due to the physical effects of the freeze-drying process, trace amounts of slowly dissolving substances may be present. This phenomenon does not affect the performance of the culture medium, and these substances will be fully utilized during cell culture. It is not caused by product contamination, so the product can be used with confidence.
- Use appropriate personal protective equipment (PPE) including gloves, lab coat, and eye protection when handling cell culture media.
- All biological materials should be treated as potentially hazardous. Follow your institution's biosafety protocols.
- Dispose of spent medium, cell culture waste, and contaminated materials in compliance with local environmental and biosafety regulations.
- Do not use after the expiration date printed on the label.

BIODOMI GMBH

Address: Schokoladenring 13, 65795 Hattersheim am Main, Germany

Document Version: 1.0 | Issue Date: March 2026